



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

319. Proposed by C. N. SCHMALL, New York City.

A man desires to purchase eggs at 5 cents, 1 cent, and $\frac{1}{2}$ cent, respectively, in such numbers that he will obtain 100 eggs for a dollar. How many solutions in rational integers?

I. Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.; J. W. CLAWSON, Ursinus College, Collegeville, Pa.; V. M. SPUNAR, Pittsburg, Pa.; and H. C. FEEMSTER, York College, York, Neb.

Let x =number at 5 cents, y =number at 1 cent, z =number at $\frac{1}{2}$ cent.
Then $x+y+z=100=5x+y+\frac{1}{2}z\dots(1, 2)$.

Eliminating z we get the indeterminate equation, $9x+y=100$.

$\therefore y=100-9x$.

This equation gives us eleven integral solutions, as follows:

$x=1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11$.

$y=91, 82, 73, 64, 55, 46, 37, 28, 19, 10, 1$.

$z=8, 16, 24, 32, 40, 48, 56, 64, 72, 80, 88$.

Also solved by S. F. Norris, S. A. Corey, B. Kramer, J. Scheffer, T. J. Fitzpatrick, and Theodore L. DeLand.

II. Solution by PROFESSOR S. F. NORRIS, Baltimore City College, Baltimore, Md., and J. E. SANDERS, Weather Bureau, Chicago, Ill.

$$\text{Average price}=1 \left\{ \begin{array}{c} 5 \\ 1 \\ \frac{1}{2} \end{array} \right\} \left| \begin{array}{c|c|c|c|c|c|c|c|c|c|c|c} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 \\ \hline 91 & 82 & 73 & 64 & 55 & 46 & 37 & 28 & 19 & 10 & 1 \\ \hline 8 & 16 & 24 & 32 & 40 & 48 & 56 & 64 & 72 & 80 & 88 \end{array} \right|.$$

NOTE. This is the method used in many arithmetics. For the process of reasoning see *e. g.*, Ray's *New Higher Arithmetic*, p. 333, subject, Alligation. ED. F.

320. Proposed by FRANCIS RUST, C. E., Pittsburg, Pa.

Solve for t , $\text{cost}=m\cos 2t$.

Solution by T. J. FITZPATRICK, Lamoni, Iowa; S. A. COREY, Hiteman, Iowa; J. E. SANDERS, Weather Bureau, Chicago, Ill.; and B. KRAMER, E. M., N. S., Pittsburg, Pa.

$$\text{cost}=m\cos 2t=m(\cos^2 t-\sin^2 t)=m(\cos^2 t-1+\cos^2 t)=2m\cos^2 t-m.$$

$$2m\cos^2 t-\text{cost}=m.$$

$$\cos^2 t - \frac{1}{2m}\text{cost}=\frac{1}{2}.$$

$$\cos^2 t - \frac{1}{2m}\text{cost} + \frac{1}{16m^2}=\frac{1}{2} + \frac{1}{16m^2}=\frac{8m^2+1}{16m^2}.$$

$$\text{cost} - \frac{1}{4m}=\pm \frac{1}{4m}\sqrt{[(8m^2+1)]}.$$